

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 51/02, 51/12, G01N 33/60, 33/84		A1	(11) International Publication Number: WO 99/04827
			(43) International Publication Date: 4 February 1999 (04.02.99)
(21) International Application Number: PCT/AU98/00582 (22) International Filing Date: 23 July 1998 (23.07.98) (30) Priority Data: PCT/AU97/00467 24 July 1997 (24.07.97) AU (71) Applicant (for all designated States except US): THE AUSTRALIAN NATIONAL UNIVERSITY [AU/AU]; Canberra, ACT 2000 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only): NAIR, Chenicheri, Hariharan [AU/AU]; 15 Bennett Place, Castle Hill, NSW 2154 (AU). SHATS, Elena, Alexandra [AU/AU]; 33 Pelham Close, Chapman, ACT 2611 (AU). BURCH, William, Martin [AU/AU]; 39 Coopernook Avenue, Gynea Bay, NSW 2227 (AU). BROWITT, Rodney, James [AU/AU]; 59 Ashburton Circuit, Kaleen, ACT 2617 (AU). SENDEN, Timothy, John [AU/AU]; 5 Bennelong Crescent, Macquarie, ACT 2614 (AU). (74) Agents: SLATTERY, John, M. et al.; Davies Collison Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: METHOD FOR DETECTION OF FIBRIN CLOTS			
(57) Abstract A method for the detection of fibrin in a source, in particular the <i>in vivo</i> detection of a fibrin in a patient, the method comprising supplying to the source or patient an amount of a detectable reagent comprising a plurality of discrete particles, each of the particles comprising a plurality of layers of carbon and being capable of binding to fibrin; and detecting the presence of the particles in the source. The particles may also comprise a detectable marker encased in said plurality of layers of carbon, the presence of said marker being capable of detection in said source.			

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

METHOD FOR DETECTION OF FIBRIN CLOTS

Technical Field

The present invention relates to a method for labelling macromolecules such as fibrin, and to methods for targeting drugs, such as antithrombotic drugs to
5 macromolecules such as fibrin in vivo. The invention also relates to a reagent for use in such a labelling or targeting method.

Background Art

10 Currently available methods of labelling macromolecules, and in particular biological macromolecules such as fibrin, may be influenced by the chemistry of the detectable marker with which the macromolecule is to be labelled. In particular, subsidiary effects resulting from the chemistry of the marker and the macromolecule being labelled may cause artifactual results. Ideally, the marker should have no influence
15 on the macromolecule to be labelled.

It is extremely important in many applications to achieve accurate and specific labelling of macromolecules such as fibrin. For example, areas of clot detection in pathological states such as deep venous thrombosis, thrombophlebitis, and lesions in the vasculature generally require accurate and specific labelling, particularly as morbidity and
20 mortality caused by intravenous clots is an increasingly prominent public health issue throughout the world.

Currently, there are several methods of detecting intravenous clots. For example, use is commonly made of two imaging agents which have similar capabilities for identifying such clots. These include fibrinogen labelled with ^{125}I and a labelled 3B6
25 monoclonal antibody. The ^{125}I labelled fibrinogen has, however, been found to be nonspecific and capable of giving false positive results.

"Technegas" is an agent whose original purpose was to perform high quality diagnostic imaging of airways of the lungs to facilitate the detection of blood clots. This was primarily used in conjunction with a blood flow agent for the differential diagnosis
30 of blood clots. The agent "Technegas" consists of a plurality of discrete particles, each

- 2 -

comprising a plurality of coating layers of carbon (varying between two and ten atoms in thickness) which completely enclose a minute crystal of ^{99m}Tc metal. The particles are stable inert hydrophobic particles having a cross-sectional dimension of between 5 and 30 nm and a thickness of about 3nm. The "Technegas" production process creates
5 millions of these particles suspended in a carrier gas of argon. In this form the particles may be inhaled directly and become deposited in the alveolar spaces of the lung. From this distribution can be created three-dimensional maps of the airways by detecting the gamma ray signal generated by the decay of the Tc from the excited to the ground state ($^{99m}\text{Tc} \rightarrow ^{99}\text{Tc}$).

10 The possibility of using the particles in liquid suspension as a "nanocolloid" suitable for biological and industrial applications has now been investigated. In particular, investigations have been made to find methods of extracting "Technegas" particles from argon gas directly into a physiological solution such as, for example, saline solution.

15

Disclosure of the Invention

It has been surprisingly found that particulate entities comprising carbon, particularly those of smaller dimensions, bind to macromolecules such as fibrin(ogen) and soluble fibrin.

20 The present invention is therefore, based on the ability of such particulate entities comprising carbon, to bind to fibrin. To do this, in vivo, the entities comprising carbon are prepared in an aqueous injectable solution (typically an isotonic solution such as 5% glucose in water) and injected intravenously. In vitro the entities may be formulated with a suitable carrier and added to a composition containing the fibrin.

25 Furthermore, it has been found that these particulate entities are advantageously able to enclose a detectable marker. This effectively quarantines the marker from its surrounding environment. Thus, the invention advantageously provides an effective means for labelling fibrin in vivo or in vitro.

According to one aspect of the invention there is provided a method for the in
30 vivo detection of fibrin in a patient, the method comprising:

- 3 -

administering to the patient an effective amount of a detectable reagent comprising a plurality of discrete particles dispersed in a veterinarily or pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant each of the particles comprising a detectable marker encased in a plurality of layers of carbon and
5 being capable of binding to fibrin; and

detecting the presence of the detectable marker in the patient.

According to another aspect of the invention there is provided a method for the detection of fibrin in a source, the method comprising:

supplying to the source a detectable reagent comprising a plurality of discrete
10 particles dispersed in a carrier, diluent, excipient and/or adjuvant, each of the particles comprising a detectable marker encased in a plurality of layers of carbon and being capable of binding to fibrin; and

detecting the presence of the detectable marker in the source.

According to a further aspect of the invention there is provided a detectable
15 reagent for use in the in vivo or in vitro detection of fibrin, the detectable reagent comprising a plurality of discrete particles dispersed in a carrier, diluent, excipient and/or adjuvant, each of the particles comprising a detectable marker encased in a plurality of layers of carbon and being capable of binding to fibrin.

The particles are distinguished from the standard "Fullerene" derivative particles in
20 that they comprise a plurality of layers of carbon. Also, their method of formation is different. Fullerenes are predominantly produced by a carbon arc process at low pressure, and the residue from the reaction is chemically and physically sorted into separate components for analysis or further processing. The subject discrete particles are preferably formed by a resistive or otherwise controlled thermal graphite heating process.
25 The co-condensation reaction in which the particles are formed in this case seems to occur in the boundary region of a stable thermal plasma that has been termed a "thermopause". Time, temperature and pressure conditions for the "thermopause" determine the size of the native metal crystal and the number of graphite layers which surround it.

In a preferred embodiment, the particles are hydrophilic. In particular, it is
30 preferred that an outer layer at least of the plurality of layers is chemically modified to

- 4 -

permit a stable chemical association of the outer layer with an aqueous medium or solution. More particularly, the outer layer preferably comprises hydrolysed graphite. Further, to ensure that the chemistry of the detectable marker is not manifested in the environment outside of the particles, the detectable marker is preferably encased in from
5 about 2 to 20 layers of graphitic carbon, more preferably about 2 to 10 layers of graphitic carbon.

Suitably, the reagent is administered to a patient in the form of an injectable composition containing a suitable carrier, for example 5% glucose in water. This carrier is preferred as it is non ionic and is generally accepted as a standard isotonic solution.
10 The total mass of the particles in one dose will generally be in the order of 100ng or less.

The detectable marker may be detected by radiometric techniques using a radionuclide, by magnetic resonance imaging, or may be optically detectable. The detectable marker may be pharmaceutically or veterinarily acceptable or otherwise (e.g. laboratory or analytically acceptable) depending on the intended use of the reagent.

15 Generally, radionuclides which are traditionally used in nuclear medicine including imaging techniques are suitable for use in this invention, and include radionuclides which emit gamma rays. Examples of suitable radionuclides are ^{111}In with a half-life of 2.8 days, ^{67}Ga , ^{68}Ga , $^{99\text{m}}\text{Tc}$. A preferred radionuclide is $^{99\text{m}}\text{Tc}$.

An example of a detectable marker suitable for use in magnetic resonance imaging
20 techniques is Gd.

Examples of markers detectable by optical techniques are colloidal compounds which exhibit a distinctive colour. An example is colloidal gold.

As discussed above, the particles used in accordance with the various aspects of the present invention differ from standard "Fullerene" derivative particles by both
25 chemistry and means of production. Accordingly, the invention also provides a method of forming a detectable reagent for use in labelling fibrin, the method comprising the steps of:

heating a carbon crucible having deposited thereon a solid form detectable marker in a sealed container to a temperature of from 2250°C to 3000°C to form particles
30 comprising a detectable marker encased in a plurality of layers of carbon and being

- 5 -

capable of binding to fibrin; and

precipitating the particles to form the detectable reagent.

The detectable marker may be deposited on the crucible in the solid form. Alternatively, a liquid solution of a detectable marker may be deposited onto the carbon
5 crucible and evaporated so as to form solid residue of the detectable marker.

Where the detectable marker is ^{99m}Tc , a liquid solution of technetium is generally used in the form of a liquid solution of sodium pertechnetate. Similarly, where the detectable label is ^{99m}Tc , the solid form of technetium is generally sodium pertechnetate.

Preferably, the crucible is heated to a temperature in the range 2250-2900°C more
10 preferably 2470-2900°C, more preferably about 2600°C to about 2700°C. In a preferred embodiment the crucible is pulse heated at from 0.1 - 5 secs, more preferably 1-3 seconds, more preferably 2.7 secs. It has been found that such pulse heating advantageously facilitates the production of smaller particles having a more tightly controllable size distribution.

15 Preferably the carbon crucible is a graphite crucible, more preferably a pure graphite crucible. The graphite may be, for example, pyrolytic graphite or porous standard graphite. However, pyrolytic graphite is preferred.

Generally the graphite crucible is heated in the presence of a substantially pure inert gas atmosphere within the sealed container. Preferably the inert gas is argon.

20 Suitable methods for preparing detectable markers such as radionuclides, encapsulated in a carbon are generally described in Australian Patent No. 589 578 (incorporated herein by cross reference). However, unlike heating the graphite crucible containing sodium pertechnetate to any temperature above at least 1900 or 2200°C under an inert atmosphere so as to produce Technegas as described in Australian Patent No. 589
25 578 it is important to only heat the graphite crucible to a temperature in the range 2250-3000°C, more preferably 2475-2950°C, more preferably 2470-2900°C, and even more preferably about 2600°C to about 2700°C. Generally pulse heating (0.5-3 secs) is used.

The particles may be precipitated using an electrostatic precipitator of the type described in AU 31778/95 sealed under Australian Patent No. 686861 on 25 May, 1998
30 (incorporated herein by cross reference), for example, and the resultant precipitate

- 6 -

then formulated with a desired carrier or adjuvant, for example at a desired concentration dependent on the end use.

The electrostatic precipitator may include a housing defining a duct through which a gas containing carbon particles passes, the duct having an inlet and an outlet; an ion source past which the gas passes to charge the particles; an electrode between the inlet and the outlet and spaced downstream from the ion source; means to establish an electric potential between the ion source and the electrode; and wherein the electrode is coated with a soluble material to which the particles are attracted so as to become deposited thereon. An electrode for the electrostatic precipitator, may include a coating of a soluble material upon which the particles are deposited by being attracted thereto. An alternative electrostatic precipitator to collect particles from a gas stream, may include a duct through which the gas passes between an inlet and an outlet; an ion source between the inlet and the outlet past which the gas passes to have the particles charged; a reservoir containing a liquid past which the gas passes; and means to establish an electric potential between the ion source and the reservoir so that particles are attracted to the liquid. Another alternative electrostatic precipitator, includes a housing defining a duct through which a gas containing carbon particles passes, the duct having an inlet and an outlet; an ion source projecting into the duct and located between the inlet and the outlet; a wall surrounding at least part of the duct between the inlet and the outlet; means in the duct to receive a liquid; means to enable the establishment of an electric potential between the ion source and the liquid; means to cause the liquid to produce droplets to be dispersed in the duct; and wherein upon the application of the electric potential, the droplets and particles are attracted to the wall. A method of collecting particles may include the steps of passing a gas stream containing the particles, through a chamber, the gas stream including an inert gas and air; passing the gas stream past an ion source within the chamber to charge the particles; and attracting the particles to an electrode by establishing an electrical potential between the ion source and the electrode. The particles may then be removed from the electrode and formulated as required.

It has been found that addition of a trace of surfactant coating to the particles may enhance their binding efficiency by up to 10 times. Preferably, the surfactant is, for

- 7 -

example, $C_{16}EO_6$. The concentration of the surfactant is preferably between 0.001% and 0.010% v/v, more preferably 0.003% and 0.006% v/v.

It has also been found that particles comprising a plurality of layers of carbon by themselves, i.e. particles which do not include via encapsulation a detectable marker, may also bind to fibrin.

Thus, the invention also provides methods for the detection of fibrin and reagents for the detection of fibrin as hereinbefore described, but which do not encase a detectable marker. In these cases, the detection of the particles to determine the presence or otherwise of fibrin, such as in the form of a fibrin clot, must be conducted by suitable means other than the abovementioned detection of the detectable marker.

According to certain embodiments of this invention wherein fibrin is detected in vitro, techniques such as dip-stick methods, test strip methods, ELISA methods, agglutination tests using latex suspensions, and the like may be suitably employed. These methods may make use of antibodies bound to the particles for example. Alternatively, in the case of latex, dip-stick methods and test strip methods antifibrin may be attached to the latex, dip stick or test strip (or other suitable substrate e.g. beads such as glass beads). Thus when such a latex particle, dip stick or test strip is dipped into or placed in contact with a source containing a fibrin: particle complex, it will bind the fibrin portion of the complex (as well as any free fibrin or other fibrin complexes in the composition). The latex particle, dip stick or test strip is then typically washed so as to remove any unattached components of the composition and whether and/or the amount of complex attached to the latex particle, dip stick or test strip determined by detecting whether the detectable label is present (and, if so, the amount present if required) on the latex particle, dip stick or test strip.

The methods of the invention are not restricted to human patients. Animal experiments have proven that this binding to fibrin occurs in deposits in deep vein thrombosis, pulmonary embolism and tumours including mammary tumours. It is envisaged that the methods of the invention may also be applicable to malignant disease such as skin cancer. The methods of the invention may be practised on mammals

- 8 -

including a bovine, human, ovine, equine, caprine, leporine, feline or canine vertebrate. Most often, however, the methods will be practised on a human patient. In this case, the composition is a pharmaceutical composition and the carrier, diluent, excipient and/or adjuvant is pharmaceutically acceptable. Where the mammal is not human, the
5 composition would be typically a veterinary composition and the carrier, diluent, excipient and/or adjuvant would be veterinarily acceptable. However, the methods of detection of fibrin in a source may be performed either in vivo or in vitro, including a reaction in a suitable laboratory apparatus such as a test tube or beaker followed by separation of any complex formed and detection for the presence of complex.

10 In this specification and claims the expression "encased" refers to total enclosure or encapsulation of the detectable marker by the carbon layers. This phenomenon may be the result of co-condensation. That is, detectable marker crystals, liquid droplets or amorphous deposits first form and then carbon is condensed around the marker to form a discrete particle. Where a marker is not required, carbon is simply prepared so as to
15 form a particle comprising a plurality of layers of carbon.

For parenteral administration, the reagent of certain embodiments of the invention may be prepared in sterile aqueous or oleaginous solution or suspension. Suitable non-toxic parenterally acceptable diluents or solvents include water, Ringer's solution, 5% glucose in water, buffered sodium or ammonium acetate solution, 1,3-butanediol, ethanol,
20 propylene glycol or polyethylene glycols in mixtures with water. Aqueous solutions or suspensions may further comprise one or more buffering agents. Suitable buffering agents include sodium acetate, sodium citrate, sodium borate or sodium tartrate, for example. Aqueous solutions for parenteral administration may also be suitable for administration orally or by inhalation.

25 The dosage form of the reagent composition will comprise typically from 0.001% to 10% by weight of the reagent. Preferably, dosage forms according to the invention will comprise from 0.1% to about 5% by weight of the reagent.

The reagent of various aspects of the invention may be prepared by means known in the art for the preparation of compositions, such as in the art of preparing veterinary
30 and pharmaceutical compositions, including blending, grinding, homogenising,

- 9 -

suspending, dissolving, emulsifying, dispersing and mixing of the reagent of the first embodiment together with the appropriate excipient(s), carrier(s), adjuvant(s) and/or diluent(s).

In the in vivo methods of the invention, the reagent may be administered orally or parenterally, e.g. by injection and by intra-arterial infusion, intravenously, rectally or by inhalation spray. A suitable method may comprise the administration of a single dose or multiple doses. If the application of more than one type of reagent is desired (e.g. there may be multiple reagents with different detectable markers) the reagents may be administered at the same time or at different times (including sequentially).

The administered dosage of the reagent(s) can vary and depends on several factors, such as the condition, age and size of the patient as well as the conditions under which the method is being performed. Dosages of reagent are preferably such that about 100ng or less of the discrete particles, in terms of the total mass of all of the particles, is administered.

For oral administration, a pharmaceutical or veterinary composition comprising the reagent may be in the form of tablets, lozenges, pills, troches, capsules, elixirs, powders, including lyophilised powders, solutions, granules, suspensions, emulsions, syrups and tinctures. Slow-release, or delayed-release forms may also be prepared, for example in the form of coated particles, multi-layer tablets or microgranules.

An inhalation spray comprising the reagent(s) may be in the form of a solution, suspension or emulsion as exemplified above. The inhalation spray may further comprise an inhalable propellant of low toxicity. Suitable propellants may include for example carbon dioxide or nitrous oxide.

Illustrative of typically used pharmaceutically or veterinarily acceptable carriers or diluents are demineralized or distilled water; isotonic glucose solution. Typically, the carrier or carriers will form from 90% to 99.9% by weight of the composition.

Suitable buffering agents include salts of boric, acetic, phosphoric, citric, malic, succinic acids and the like, for example sodium citrate, sodium bicarbonate, sodium acetate and sodium phosphate. Additionally or alternatively, the free acids may be used, together with an alkali such as sodium hydroxide, sodium carbonate, sodium bicarbonate,

- 10 -

potassium hydroxide, potassium carbonate or potassium bicarbonate. Typically, the buffering agent or agents will form from 0.1% to 20% by weight of the composition.

The following expressions are used in the Examples.

a) **FullerTag**: (Registered Trademark) A nanocolloid suspension (in saline or distilled water) of particles comprising 2-10 atomic layers of carbon encasing technetium crystal such that it forms a stable inert hydrophobic or hydrophilic particle whose overall dimension is between about 5 and about 30nm in cross-section and about 3nm in thickness. b) **ThromboTrace**: *FullerTag* with surfactant. In the Examples, the surfactant is $C_{16}EO_6$.

10 The specific affinity of the multi-layered carbon particles for fibrin, whether containing or not containing a detectable marker, suggests that the invention may also be relevant to systems for "site-specific" drug delivery. In particular, it is envisaged that the affinity of the particles of the invention to fibrin may be useful in targeting, for example, antithrombotic or anti-cancer drugs to fibrin sites in order to provide more direct and effective treatment.

15 Accordingly, the invention also provides a method of targeting a drug to a fibrin site in vivo, the method comprising administering to a patient an effective amount of a reagent comprising a plurality of discrete particles dispersed in a veterinarily or pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant, each of said particles comprising a plurality of layers of carbon and being capable of binding to fibrin, and at least some of said particles having coupled thereto a drug to be targeted to the fibrin site.

25 Accordingly there is also provided a method of treatment of a fibrin related disease comprising administering to a patient an effective amount of a reagent comprising a plurality of discrete particles dispersed in a veterinarily or pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant, each of said particles comprising a plurality of layers of carbon and being capable of binding to fibrin, and at least some of said particles having coupled thereto a drug to be targeted to the fibrin site.

30 As mentioned above, the drug to be targeted may include any drug, for example anti-thrombotic or anti-cancer drugs. Furthermore, these drugs may be coupled to the

- 11 -

particles by any suitable means including coupling methods known in the art. The particles may also encase a detectable marker as previously described. In this instance, the particles may not only target the coupled drugs to a fibrin site, but may also be used to detect the location of the fibrin site in vivo.

5 Following from this, the invention also provides a reagent for use in targeting a drug to a fibrin site in vivo, the reagent comprising a plurality of discrete particles dispersed in a pharmaceutically or veterinarily acceptable carrier, diluent, excipient and/or adjuvant, each of the particles comprising a plurality of layers of carbon and being capable of binding to fibrin, wherein at least some of the particles have coupled thereto
10 a drug to be targeted to the fibrin site.

Brief Description of the Drawings

Embodiments of the invention will now be described in more detail with reference to the accompanying drawings in which:

15 Figure 1 is an autoradiograph of a fibrin clot in accordance with Example 12;
 Figure 2 is a schematic representation of the artificial circuit described in Example
13;

 Figure 3 shows the accumulation of radioactivity in the intramural clot in the artificial circuit of Example 13;

20 Figure 4 shows the transmission electron micrography described in Example 14;
 Figure 5 shows imaging of clots in a rabbit ear vein described in Example 17;
 Figure 6 shows a clot caused by intramuscular injection of anaesthetic described
in Example 18;

 Figure 7 shows the imaging of a clot in a cat femoral vein described in Example
25 19;

 Figure 8 shows the imaging of a clot in rabbit pulmonary vasculature described
in Example 20;

 Figure 9 shows the autoradiography of rabbit lungs with pulmonary emboli also
described in Example 21;

30 Figure 10 shows the labelling of a small tumour in the ear of the rabbit as

- 12 -

described in Example 23;

Figure 11 shows the labelling of a mammary tumour in a Fischer strain rat as described in Example 23; and

Figure 12 shows particles with colloidal gold.

5

Best Modes and Other Modes for Carrying the Invention

It is envisaged that the specific binding of particles according to the invention may be useful for:

- a) imaging of clots in blood vessels in patients with deep vein thrombosis
- 10 (DVT), patients with atheromatous occlusions in the vasculature and embolisms;
- b) the development of target specific drugs for the dissolution of blood clots;
- c) the use of technology to develop a bedside measurement of fibrin(ogen) and soluble fibrin;
- d) the detection of early tumours; and
- 15 e) research tools to study clots and diseases like atherosclerosis, Deep Vein Thrombosis and any other fibrin related disease.

The mechanism for formation of the carbon particles depends on a co-condensation of the native metal crystals with the C_3 gas that is present above carbon at high temperatures. The reaction is carried out in an inert gas atmosphere, typically argon.

20 Thus, the final size of the nano-encapsulates can be modified by varying the rate and duration of the heating cycle. At present, repeat short pulses of heat for 2.7 sec to temperatures as high as from 2600°C to 2700°C will produce particles that go to the lower limit of resolution of electron microscopes, namely 5nm. Pyrolytic or glassy-carbon provides a better yield of activity of Tc from the crucible than the more porous

25 standard graphite. This appears to be due to the non-absorption of the ions into the graphite matrix. Four lots of 2.7sec heating pulses to from 2600°C to 2700°C in pyrolytic or glassy-carbon gives rise to a much smaller spectrum of particle size distribution than the original heating cycle of 2500°C for 15 sec. The particles were precipitated on an electrode using a precipitator as described in Australian Patent

30 Application No. 31778/95, sealed under Australian Patent No. 686861 on 25 May, 1998

- 13 -

and formulated in saline. The reagent composition may be then injected into a patient and detected in vivo so as to locate fibrin deposits.

Typically the starting material is Tc-99m as standard sodium pertechnetate for injection. The Micro-Aerosol Generator (MAGen - USA name), or particle generator is essentially a miniature high temperature furnace in which the heating element is also the source of graphite and/or carbon vapour which ultimately coats the Technetium metal. The heating element is made from 100% pure spectroscopic graphite whose electrical and mechanical specifications match the requirements of the machine. It is a 6mm square section rod 50mm long, machined to form a crucible in the centre section such that it can hold a liquid volume of 0.14mL. This hollowed and thinned section also provides the high resistive portion of the rod which becomes the hottest section when electric current is passed through it. The rod is held under spring tension between two high current electrodes. The entire assembly is mounted as a drawer section sliding into the lower chamber of a 6L vessel, and electrically powered from an automatic process-controller. The crucible is first filled with liquid Sodium Pertechnetate in normal physiological saline, which in most instances contains enough activity [260-370MBq or 7-10mCi] for a single patient administration. The drawer section is then closed and the automatic process takes over, gently blowing pure argon gas over the top of the crucible while warming it to 70°C. This is known as the "simmer" cycle, and takes 6 minutes, during which time the liquid in the crucible dries out and the whole chamber is purged with pure argon, replacing all the original air and water vapour. A process controller alerts the user that the machine is ready to produce the particles. At the conclusion of this preparation phase, the machine is activated via the control panel "start" button, and the crucible temperature rises to 2550°C by resistively heating it with about 2.5kW of power within 0.75s (four lots of 2.7sec heating pulses to 2700°C in pyrolytic carbon gives rise to a much smaller spectrum of particle size distribution than the original heating cycle of 2550°C for 15 sec) and holds that value within 50°C for 15s through a feedback servo from an optical sensor before switching off. This fills the 6L chamber with gas comprising the particles which encapsulate Tc-99m. The particles are collected by passing the gas stream containing the particles, through a chamber, the gas stream

- 14 -

including an inert gas and air; passing the gas stream past an ion source within the chamber to charge the particles; attracting the particles to an electrode by establishing an electrical potential between the ion source and the electrode. The particles are then removed from the electrode and formulated with a carrier as required.

5 The present invention will now be described with reference to the following examples which should not be construed as limiting on the scope thereof.

Example 1

 Particles were prepared in accordance with the methods of Australian Patent No. 589 578 except the graphite crucible was subjected to four lots of 2.7sec heating pulses to 2700°C in pyrolytic or glassy-carbon. The disclosure of Australian Patent No. 589 578 is hereby incorporated by reference.

Example 2

15 Particles were formulated into normal saline by precipitating the particles formed as per Example 1 using precipitation apparatus and methods of Australian Patent No. 686861 the disclosure of which is hereby incorporated by reference, and then mixing the desired amount of the particles with normal saline.

Example 3

20 Plasma was mixed with a small aliquot of particles formulated into saline in accordance with Example 2 and clotted with 1 U/mL thrombin. After incubation at room temperature for 1 hour the plasma clot was crushed using a wooden spatula. A gamma counter was used to count radioactivity. Radioactivity was confined to the fibrin with less
25 than 3% of the original activity resident in the exudate.

- 15 -

Example 4

A similar experiment using purified fibrinogen was performed to exclude binding to other plasma proteins. Again it was shown that only about 3% of the radioactivity was resident in the exudate.

- 5 Both these experiments show that these particles have specific binding capacity for fibrin.

Example 5

- 10 To ensure that such labelling could not be transferred or leached from clots to whole blood or plasma, plasma clots were incubated in whole blood or plasma at 37°C for 3 hours. The radioactivity was confined to the clot with very little escaping to the surrounding blood or plasma (0.7% radioactivity in blood and plasma, respectively).

Example 6

- 15 Unlabelled plasma clots and clots developed from purified fibrinogen solution in glass tubes respectively were gently permeated with particles formulated into saline in accordance with Example 2. The initial washout of serum was devoid of radioactivity as expected. Further measurement showed that most of the particles were specifically bound to fibrin. These clots were removed from the glass tubes and crushed and carefully
20 washed three times in saline and counted in a gamma-counter. Radioactivity was confined to the clot and barely detectable in the exudate.

Example 7

- 25 Tests of the efficacy of the fibrin label were performed in vivo, using the rabbit, which is the standard model for vascular research. This experiment was performed on multiple occasions on different animals. One ear was given a small needle stick injury and particles formulated into saline in accordance with Example 2 in 1mL volume was injected in the other ear. The rabbit was anaesthetised for the whole procedure and laid on the detector surface of a gamma camera (in this instance a GE model 400T). In the
30 second animal there was a further small vascular injury and bruise in the ear on the side

- 16 -

of the injection.

The injury on the ear opposite to the injection site showed an increased uptake on the "first pass" of the blood flow through the body. In the second rabbit, the injury proximal to the injection site labelled much more strongly than that in the other ear, but
5 then washed out over a period of twenty minutes. Dynamic analysis showed that the initial uptake and the wash-out followed by a renewed uptake slowly with time, occurred in both animals at about the same rate.

Examples 8-16 and 22-25 relate to in vitro studies while Examples 17-21 relate to in vivo studies.

10

Example 8

Binding of *ThromboTrace* to fibrin

Protocol: Fibrin network developed from 1 mL pooled human plasma obtained from healthy donors (by adding 1U/mL bovine thrombin(Parke Davis) and 25mM CaCl_2
15 (final concentration)) was incubated with normal saline (NS) (4mL) containing *ThromboTrace* (TT) (100 μL) at 37°C for 2 hours using an elliptical rotator. A mixture of normal saline and *ThromboTrace* without clot was used as control. Clots were then crushed with a wooden spatulum and counted using a gamma counter. Nanocolloid preparation of $^{99}\text{Tc-Sb}_2\text{S}_3$ (antimony sulphide) used for bone imaging was studied as a
20 control.

Results:

50 μ l NS + <i>TT</i> (initial count)	99500 cpm	50 μ l NS + $^{99}\text{Tc-Sb}_2\text{S}_3$ (initial count)	397800cpm
50 μ l NS + <i>TT</i> (post-incubation)	87450cpm	50 μ l NS + $^{99}\text{Tc-Sb}_2\text{S}_3$ (post-incubation)	415000cpm
50 μ l NS + <i>TT</i> + clot (post incubation)	72870cpm	50 μ l NS + $^{99}\text{Tc-Sb}_2\text{S}_3$ + clot (post-incubation)	408700cpm
Crushed clot on spatulum	478720cpm	Crushed clot on spatulum	9430cpm
After extensive wash in NS	457436cpm	after 3 washes in NS	7790
		after 6 washes in NS	3150

10

Conclusion: There is a significant decrease of the label (17%) in the incubation medium with *ThromboTrace* which accounts for the high count observed in the crushed fibrin clot. There is no significant changes in the activity of incubation medium with nanocolloid and there was no significant binding of $^{99}\text{Tc-Sb}_2\text{S}_3$ to fibrin, after 6 washes most of the activity washes away. This observation suggests a high specificity of *ThromboTrace* for fibrin.

15

Example 9

Investigation of the strength of *ThromboTrace* binding to fibrin.

20

Protocol: *ThromboTrace* (*TT*) (50 μ l) was added to pooled plasma obtained from the blood of healthy donors (5mL). Fibrin networks were developed from 1 mL of this labelled plasma (by adding 1U/mL bovine thrombin (Parke Davis) and 25mM CaCl_2 (final concentration)) and allowed to incubate at 37°C for 30 min. Clots were then crushed with wooden spatulum, extensively washed with normal saline and counted using a gamma counter. Nanocolloid preparation of $^{99}\text{Tc-Sb}_2\text{S}_3$ (antimony sulphide) used for bone imaging was studied as a control.

25

Results:

- 18 -

	50µl plasma + <i>TT</i> (initial count)	58700cpm	50µl plasma + $^{99}\text{Tc-Sb}_2\text{S}_3$ (initial count)	100195cpm
	50µl expelled serum (after clot crushed)	1600cpm	50µl expelled serum (after clot crushed)	100500cpm
5	Crushed clot on spatulum	1026700cpm	Crushed clot on spatulum	3650cpm
10	Crushed clot on spatulum after extensive wash in normal saline	1005850cpm	Crushed clot on spatulum after extensive wash in normal saline	1800cpm

Conclusion: Fibrin clot retained 97% of the initial radioactivity of *TT* solution which was tightly bound. There was no binding of nanocolloid $^{99}\text{Tc-Sb}_2\text{S}_3$ to fibrin.

Example 10

15 Study of the washout of *ThromboTrace* from clot.

Protocol: *ThromboTrace* (*TT*) (50µl) was added to pooled plasma obtained from the blood of healthy donors (5mL). Fibrin networks were developed from 1 mL of this labelled plasma (by adding 1U/mL bovine thrombin (Parke Davis) and 25mM CaCl_2 (final concentration)) and allowed to incubate at 37°C for 30 min. The clots were
20 incubated 37°C for 3 hours using an elliptical rotator with 3mL unlabelled whole blood or 3mL unlabelled plasma. The radioactivity of incubation media was measured using a gamma counter.

Results:

	50µl plasma + <i>TT</i> (initial count)	58700cpm
25	50µl blood incubated with labelled clot	380cpm
	50µl plasma incubated with labelled clot	450cpm
	Crushed clot on spatulum	1026700cpm

Conclusion: *ThromboTrace* was tightly bound to fibrin, washout of label into surrounding blood or plasma was less than 1%.

- 19 -

Example 11**Study of the binding of *ThromboTrace* to fibrin under perfusion.**

Protocol: Fibrin network was developed in pre-etched glass tube from 1 mL purified fibrinogen solution (2.5mg/mL) by adding 1U/mL bovine thrombin (Parke Davis) and 25mM CaCl_2 (final concentration). The network was gently perfused with 1.5mL normal saline containing *ThromboTrace* (TT) (50 μl). The radioactivity of expelled perfusate was measured using gamma counter. Clots were then crushed with a wooden spatulum washed in normal saline and counted using a gamma counter.

Results:

10	50 $\mu\text{ NS + TT (initial count)$	47500cpm
	50 $\mu\text{l expelled perfusate}$	3200cpm
	Crushed clot on spatulum washed with NS	991200cpm

Conclusion: After the perfusion of normal saline containing *ThromboTrace* through fibrin network only 7% of initial radioactivity was detected in the perfusate while crushed clot retained considerable radioactivity. This observation indicates a high affinity of *ThromboTrace* to purified fibrin. This experiment was repeated with plasma clots and similar results were observed.

Example 12**Study of the specific binding of *ThromboTrace* to fibrin by autoradiography.**

Protocol: Fibrin network was developed in pre-etched glass tube from 1 mL purified fibrinogen solution (2.5mg/mL) by adding 1U/mL bovine thrombin (Parke Davis) and 25mM CaCl_2 (final concentration). The network was gently perfused with 1.5mL normal saline containing *ThromboTrace* (TT) (50 μl). The tube was autoradiographed with a gamma camera (Fig. 1).

Conclusion: Clots layered with *ThromboTrace* and autoradiographed with a gamma camera revealed extensive binding of material within the first cm of perfusion of the clot.

- 20 -

Example 13

Study of the binding of *ThromboTrace* to fibrin in dynamic situation using an artificial circuit.

Protocol: The artificial circuit is an adaptation of the Chandler Loop attempting
5 to simulate the vascular environment of the body. This experimental set up allows the
monitoring of the imaging of clots with the control of the parameters of flow and pressure.
Fibrin network was developed from pooled human plasma obtained from healthy donors
(by adding 1U/mL bovine thrombin (Parke Davis) and 25mM CaCl_2 (final concentration))
in a specially designed glass cell with a depressed well to house the clot. This design
10 simulated an intramural clot with its surface exposed to the flowing medium. The clot
was allowed to incubate at 37°C for 30 min. The cell was connected with silicone tubing
to the circuit comprised of a peristaltic pump and manometer (**Fig. 2**).

The circuit was filled with 50mL normal saline containing 10% pooled human
plasma and 1 mL *TT* (2mCi/mL). The cell with clot was tilted so that the circulation
15 mixture was not allowed to accumulate on top of the clot and was only trickling along the
surface of the clot. A separate hanging clot was developed within a loop of copper wire
and introduced into the circuit so that the total surface of the clot was exposed to the
circulating media. Images were dynamically recorded using a gamma camera. Data
acquisition time was 30 min. After that the circulation mixture was removed, the circuit
20 was washed with normal saline and the image of the clot was acquired for another 10
minutes.

Results:

Fig. 3 shows the image of the clot after 30 min of dynamic labelling with
ThromboTrace and after the silicone tubes and the clot were washed with normal saline.
25 The bright spot in the circuit above the clot is that of the hanging clot formed within a
small copper wire loop and simulating small clots found in the body.

Conclusion: It is obvious that both clots have been labelled with *ThromboTrace*
and show up clearly.

- 21 -

Example 14**Study of the binding of *ThomboTrace* to fibrin using Transmission Electron Microscopy.**

Protocol: Fibrin networks were developed from purified fibrinogen solution or plasma containing *TT* (by adding 1U/mL bovine thrombin (Parke Davis) and 10mM CaCl_2 (final concentration)). They were primary fixed in 2.5% glutaraldehyde and postfixed in 2% osmium tetroxide. They were then stained en block with 2% uranyl acetate, washed in distilled water, dehydrated and embedded in Spurr resin. The samples were cut on a Reichert-Jung Ultracut, stained with Reinold's lead citrate stain and micrographs taken on Philips CM 10 TEM. Negative stained fibrin clots with *TT* were prepared on carbon grids preliminarily treated with glow discharge. The grids were washed in distilled water, dried and stained in 1% sodium phosphotungstate. They were then briefly washed and dried. Subsequently, micrographs were taken on a Philips 301 TEM.

Results and Conclusions: Fig. 4 shows transmission electron micrography which reveals particles adhering to fibrin along fibres and specifically bound to dark bands of negatively stained fibrin thus denoting a specific binding site for particles.

Example 15**Study of the binding of *ThromboTrace* to fibrin monolayer using microwell technique.**

Protocol: 96 well PVC plates were coated with one of the following: PBS, solution of bovine serum albumin (BSA)/PBS. D-dimer, soluble fibrin, fibrin. The concentration of proteins directly coating well was 10 $\mu\text{g/mL}$. If proteins were raised from the surface by using 4D2/182 monoclonal antibody specific to fibrin(ogen) (AGEN) (10 $\mu\text{g/mL}$), the concentration of proteins was 100 $\mu\text{g/mL}$. Wells were washed 3 times with PBS/BSA buffer in between application of proteins or *TT*. Plates were incubated with *TT* at 37°C for 30 min. Plates were washed with PBS/BSA buffer and each well was counted using a gamma-counter.

Results:

- 22 -

Protein	% binding of <i>TT</i> compared to BSA-treated wells
Fibrinogen	38
Plasma + thrombin	177
D-dimer	0
Fibrin	179
Soluble Fibrin	189

Conclusions: *ThromboTrace* binds with high affinity to fibrin monolayer and soluble fibrin. This affinity is not influenced by albumin or fibrinogen. In separate experiments it has been demonstrated that increasing the amount of fibrinogen does not inhibit the binding of *TT* to fibrin. *ThromboTrace* does not bind to D-dimer or albumin.

Example 16

Study of the effect of surfactant (C₁₆E₀₆) on the binding of *ThromboTrace* to fibrin.

Protocol: The studies have been performed using multiwell technique (Example 15). *TT* solution was prepared with different concentration of surfactant.

Results:

Concentration of the surfactant (%)	% binding of <i>TT</i> compared to concentration of the surfactant
0.0015	140
0.003	258
0.00625	194
0.0125	168
0.025	133
0.05	139

Conclusions: Binding is enhanced with C₁₆E₀₆. Maximal binding is observed at the concentration of 0.003%.

The *in vivo* experiments described below were carried out on rabbits and a cat

- 23 -

adhering to standard anaesthetising procedure according to the guidelines provided by the Australian National Health and Medical Research Council. All animals were injected with 1 ml. *ThromboTrace* (2mCl/mL). Acquisition time for all images was 60 min.

5

Example 17

Imaging of clots in rabbit ear vein.

Two needle stick injuries with 20G needles were made in the right ear and one in the contralateral ear. *ThromboTrace* was injected into the ear vein of the right ear. As shown in Figure 5 both right ear and contralateral ear injuries have been labelled with

10 *ThromboTrace*.

Example 18

Clot caused by intramuscular injection of anaesthetic.

Fig 6 shows a clot caused by intramuscular injection of anaesthetic in the left thigh. *ThromboTrace* was injected into the ear vein. This photograph is representative of imaging done both one hour after the injection and 24 hours post injury.

15

Example 19

Imaging of clot in cat femoral vein.

20 Needlestick injury was made in the femoral vein of a cat. *ThromboTrace* was introduced near the site of injury after one hours. The labelled clot is clearly visible in the Figure 7. Further along the site of injury is a set of four bright spots. It is suspected that these were clots formed by previous injury to the animal. It was not possible to confirm this finding.

25

- 24 -

Example 20

Imaging of clot in rabbit pulmonary vasculature

Human blood clot was developed around a cotton thread in a polyurethane tube (diameter=3mm) with 1U/mL thrombin and 10mM CaCl_2 . After washing with normal
5 saline a 1mm length clot was introduced through a 16 or 18G catheter into the jugular vein of a rabbit. After 10 min *ThromboTrace* was introduced through the catheter or through the central ear artery. The results are shown in Figure 8. Classic wedging denoting depletion of vascular supply was seen in all cases.

10

Example 21

Autoradiography of rabbit lungs with pulmonary emboli

Lungs were removed from the rib cage and autoradiographed to match images of clots obtained *in vivo*. The introduced human clot with cotton thread was then identified in the lung on post mortem and confirmed by two other witnesses and agreed that images
15 obtained during the *in vivo* acquisition were in fact the introduced human blood clot. The results are shown in Figure 9.

Example 22

Specificity of particles for equine fibrin

20 *In vitro* experiments using plasma clots developed from equine blood also shows similar specificity of particles for equine fibrin. Studies utilised methods as described in Example 9. No significant difference was found between the binding characteristics of *TT* to horse fibrin and human fibrin.

Example 23

25

Binding to tumours

Fig. 10 shows the labelling of a small tumour on the ear of a rabbit. The experiment was performed as described in Example 7. Studies show that tumours on rabbit ear can be labelled using *ThromboTrace*, and the particles may thus be used in labelling tumours and cancers which have a fibrin sheath. Further conclusive evidence of this can be seen with
30 reference to Fig. 11 which shows the labelling of a mammary tumour in a Fischer strain

- 25 -

rat. In this case, cancer cells were injected subcutaneously and allowed to grow for a period of seven days. ThromboTrace was injected through the tail vein. The rat on the left of Fig. 11 is a control animal without cancer.

Example 24

5

Bone Scan

Bone breakages are followed by a deposition of fibrin. In particular, hairline fractures are often difficult to distinguish and diagnose. The propensity of the particles to bind to fibrin provides another application as a label for bone fractures and therefore a frontline bone scan.

10

Example 25

Binding to soluble fibrin

High affinity binding of *ThromboTrace* to soluble fibrin can be used as a basis for a diagnostic test for soluble fibrin which is reportedly a marker for cardiovascular disease and cancer among other diseases, e.g. Bredbacka *et al.* (1994 a & 1994 b), Ginsberg *et al.*,
15 (1995), Ginsberg *et al.* (1996), Iversen *et al.* (1995), Nakagawa *et al.* (1994), Shaukat *et al.* (1995). Experimental data set out in Example 15 demonstrates the binding of *ThromboTrace* to soluble fibrin.

20

Example 26

Diagnostic kit for soluble fibrin

Soluble fibrin bound particles are isolated and a suitable monoclonal antibody specific for soluble fibrin is attached to the soluble fibrin. An ELISA assay is then developed to specifically detect the quantity of the soluble fibrin in plasma.

25

Colloidal gold may be incorporated into these particles in order to strengthen the integrity of particles and increase the mass, thus increasing the sensitivity of the immunogold assay.

The affinity of the particles to soluble fibrin can then be used to detect the quantity of soluble fibrin again by utilising the immunogold type technique. Alternatively, it may
30 be possible that the aggregation of particles caused when they bind to soluble fibrin may

- 26 -

be used as the method of detection as these particles may exhibit a subtle colour change because of the aggregation or difference in light scattering.

Example 27

5

Targeting Specific Drugs

A novel use for the particles is the coupling of anticoagulant drugs or clot specific drugs to the carbon shell. This can be established by the functionalisation of the carbon by amination or hydroxylation. These procedures will be accomplished using a radiofrequency plasma field. Once functionalised these particles will then be coupled to
10 drug molecules using commercially available kits. In fact one anticoagulant already available is biotinylated heparin which can then be coupled to the carbon shell.

Coupling to a drug is followed by Enzyme Immunoassay to ensure binding to fibrin is not disrupted. These tests also show the effectiveness of these drugs when bound to the carbon. These coupled particles are then injected into an animal. The first
15 injections are with radioactive technitium based particles which enables detection and further describes the kinetics of dissolution of the clots. If particles are coupled with tissue plasminogen activator for example, this indicates a rapid dissolution of the clot. If particles are heparin coupled these experiments will provide ample evidence that the heparin is in fact delivered to the site it is required. This is one major disadvantage of
20 heparin usage at the moment.

Industrial Applicability

It should be clear that the methods of detection of this invention will find wide use in the medical and veterinary fields.

25

The foregoing describes only some embodiments of the present invention and modifications obvious to those skilled in the art can be made thereto without departing from the scope of the invention.

References

- Bredbacka *et al.* Soluble fibrin: a predictor for the development and outcome of multiple organ failure. *Am J Hematol* 46:289-294, 1994 (a)
- 5 Bredbacka *et al.* Bredbacka *et al.* Soluble fibrin and D-dimer as detectors of hypercoagulability in patients with isolated brain trauma. *J Neurosurg Anesthesiol* 6:75-82, 1994 (b)
- Ginsberg *et al.* Evaluation of a soluble fibrin assay in patients with suspected deep vein thrombosis. *Thromb Haemost* 75(3):833-836, 1995.
- 10 Ginsberg *et al.* Evaluation of a soluble fibrin assay in patients with suspected pulmonary embolism. *Thromb Haemost* 75(4):551-554, 1996
- Iversen *et al.* Soluble fibrin in plasma before and after surgery for benign and malignant colorectal disease. *Thromb Res* 79:471-481, 1995
- Nakagawa *et al.* Plasma levels of soluble fibrin in patients with malignancy-associated
- 15 disseminated intravascular coagulation. *Blood Coagul Fibrinolysis* 5:725-730, 1994
- Shaukat *et al.* Soluble fibrin: a marker of the development of thrombosis in coronary heart disease. *Thromb Haemost* 73(6):1141 (922), 1995.

- 28 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for the in vivo detection of fibrin in a patient, said method comprising:
 - administering to said patient an effective amount of a detectable reagent
 - 5 comprising a plurality of discrete particles dispersed in a pharmaceutically or veterinarily acceptable carrier, diluent, excipient and/or adjuvant, each of said particles comprising a detectable marker encased in a plurality of layers of carbon and being capable of binding to fibrin; and
 - detecting the presence of said detectable marker in said patient.
- 10 2. A method for the detection of fibrin in a source, said method comprising:
 - supplying to said source a detectable reagent comprising a plurality of discrete particles dispersed in a carrier, diluent, excipient and/or adjuvant, each of said particles comprising a detectable marker encased in a plurality of layers of carbon and being
 - capable of binding to fibrin; and
 - 15 detecting the presence of said detectable marker in said source.
3. A method according to claim 1 or 2, wherein each of said plurality of particles is hydrophilic and comprises said detectable marker encased in from 2 to 10 layers of graphitic carbon.
4. A method according to claim 1 or 2, wherein the carrier is an aqueous medium
- 20 or solution.
5. A method according to claim 4, wherein the aqueous medium or solution is 5% glucose in water.
6. A method according to claim 1 or 2, wherein each of said discrete particles has a cross-section of from about 5nm to about 30nm.
- 25 7. A method according to claim 1 or 2, wherein said reagent is administered in an amount such that the dose comprises about 100 ng or less of said particles..
8. A method according to claim 1 or 2, wherein said detectable marker is detectable by radiochemical techniques, magnetic resonance imaging or is optically detectable.
- 30 9. A method according to claim 1 or 2, wherein said detectable marker is selected

- 29 -

from the group consisting of radionuclides which emit gamma rays; Gd or Au.

10. A method according to claim 9, wherein said detectable marker is ^{99m}Tc .

11. A detectable reagent for use in the in vivo or in vitro detection of fibrin, said detectable reagent comprising a plurality of discrete particles dispersed in a carrier, 5 diluent, excipient and/or adjuvant, each of said particles comprising a detectable marker encased in a plurality of layers of carbon and being capable of binding to fibrin.

12. A detectable reagent according to claim 11, wherein each of said plurality of particles comprises a detectable marker encased in from 2 to 10 layers of graphitic carbon, at least an outer layer of said layers being chemically modified to permit a 10 stable chemical association of the layer with aqueous medium or solution.

13. A detectable reagent according to claim 11, wherein the outer layer comprises hydrolysed graphite.

14. A detectable reagent according to claim 11, wherein the carrier is an aqueous medium or solution.

15 15. A detectable reagent according to claim 14, wherein the aqueous medium or solution is 5% glucose in water.

16. A detectable reagent according to claim 11, wherein each of said particles has a cross-section of from about 5nm to about 30nm.

17. A detectable reagent according to claim 11, wherein the detectable marker is 20 detectable by radiometric techniques, magnetic resonance imaging or is optically detectable.

18. A detectable reagent according to claim 11, wherein said detectable marker is selected from the group consisting of radionuclides which emit gamma rays; Gd or Au.

19. A detectable reagent according to claim 18, wherein said detectable marker is 25 ^{99m}Tc .

20. A method of targeting a drug to a fibrin site in vivo, the method comprising administering to a patient an effective amount of a reagent comprising a plurality of discrete particles dispersed in a veterinarily or pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant, each of said particles comprising a plurality of layers 30 of carbon and being capable of binding to fibrin, and at least some of said particles having

- 30 -

coupled thereto a drug to be targeted to the fibrin site.

21. A method according to claim 20, wherein said drug to be targeted is an anti-thrombotic or anti-cancer drug.

22. A method according to claim 20, wherein each of said particles comprises a
5 detectable marker enclosed in said plurality of layers of carbon.

*SPECIFIC BINDING OF THROMBO TRACE TO
SURFACE OF A CLOT AS SEEN BY AUTORADIOGRAPHY.*





FIG. 1

*ARTIFICIAL CIRCUIT**FIG. 2*

*THE ACCUMULATION OF RADIOACTIVITY IN THE
INTRAMURAL CLOT IN AN ARTIFICIAL CIRCUIT.*




FIG. 3

*TRANSMISSION ELECTRON MICROGRAPHY OF NEGATIVELY
STAINED FIBRIN WITH THROMBO TRACE.*




FIG. 4

IMAGING OF CLOTS IN RABBIT EAR VEIN.




FIG. 5

*IMAGING OF A CLOT IN RABBIT CAUSED BY
INTRAMUSCULAR INJECTION*




FIG. 6

IMAGING OF A CLOT IN THE CAT FEMORAL VEIN.




FIG. 7

*IMAGING OF A CLOT IN RABBIT PULMINARY
VASCULATURE.*




FIG. 8

*AUTORADIOGRAPHY OF RABBIT LUNGS WITH
PULMINARY EMBOLI.*




FIG. 9

*LABELLING OF A SMALL TUMOUR ON THE EAR OF
A RABBIT*





FIG. 10

FIG 11







FIG. 12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU 98/00582

A. CLASSIFICATION OF SUBJECT MATTER				
Int Cl ⁶ : A61K 51/02, 51/12 G01N 33/60, 33/84				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED				
Minimum documentation searched (classification system followed by classification symbols) IPC : A61K and keywords, G01N 33/- and keywords				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) WPAT (DERWENT) : (Carbon or Graphit:) and Fibrin: and (A61K or G01N 33/-) CHEMICAL ABSTRACTS : (Carbon or Graphit?) and (layer? or mesh? or ?Encapsul?:) and Fibrin?				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X Y	WO 93/15768 A (NYCOMED SALUTAR INC) 19 August 1993 page 6 line 8 to page 7 line 6, page 18 line 4 to page 19 line 22 page 29 lines 1 to 19	11-19 1-22		
P,Y	Journal of Biomedical Materials Research, Volume 37, No. 1, issued November 1997 (John Wiley & Sons, Inc), H. Nygren and C. Eriksson, "The Initial Reactions of Graphite and Gold with Blood", pages 130-136 whole document	1-22		
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex </div>				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top;"> <p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </td> <td style="width: 50%; vertical-align: top;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p> </td> </tr> </table>			<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>			
Date of the actual completion of the international search 25 September 1998		Date of mailing of the international search report 30/sep/1998		
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200 WODEN ACT 2606 AUSTRALIA Facsimile No.: (02) 6285 3929		Authorized officer  ANDREW ACHILLEOS Telephone No.: (02) 6283 2280		

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 98/00582

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Biomaterials, Volume 8, No. 1, issued January 1987, (Butterworth & Co Ltd), E. Chigmier et al, "Haemocompatibility and Biological Course of Carbonaceous Composites for Cardiovascular Devices", pages 18-23 whole document	1-22
Y	Proc of the Thirty-Eighth Annual Meeting, Electron Microsc. Society Am., issued 1980, (EMSA), M.L. Rudee et al, "The interaction of Blood with Amorphous Carbon", pages 584-5 whole document	1-22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU 98/00582

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report				Patent Family Member			
WO	93/15768	AU	35068/93	EP	625055	GB	9203037
		US	5688486				

END OF ANNEX

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)